

Driclor

SOLUTION

TEXT FOR CONSUMERS

EN / Feeling nervous, as well as high temperature, humidity or physical exertion, may lead to increased sweating. Driclor creates a gel seal in the outlet of the sweat gland, preventing sweat from escaping at the application site. Sweat is excreted from the body in a natural way (e.g. through the body surface, kidneys). After a few weeks of use, excessive sweating is reduced and the patient has a feeling of dryness and well-being. **Adults and adolescents aged 12+.**

DOSAGE AND USE

For external use only.

Driclor should only be used on the armpits, palms and soles. Driclor should only be used on thoroughly dried skin, especially after a bath, otherwise it may cause skin irritation or burning.




In the initial stage, Driclor should be used once a day, as the last activity before sleep.

Wash the treated areas the next morning. Do not use Driclor during the day.

Once sweating begins to subside during the day, you will be able to reduce the frequency of using Driclor to 1-2 times a week or less often.

INFORMATION ON PREPARATION OF THE DRICLOR PACKAGE

1. Place the bottle on a flat surface, carefully unscrew the cap and set it aside.
2. Insert the ball applicator into the neck of the bottle and push it gently but firmly.
3. Seal the cap well after use. Store the bottle with the cap facing up.



Do not use:

— If you are allergic to any of the ingredients of the product.

WARNINGS AND PRECAUTIONS

- Do not use Driclor on injured or irritated skin.
- Do not use hair removal creams and do not shave your armpits for at least 12 hours before and after using Driclor.
- Do not use Driclor around the eyes, nose and mouth.
- Driclor is highly flammable. When using the product, do not smoke and stay away from naked flame.
- Driclor can cause stains on clothes, so dry the treated area thoroughly before dressing.
- Avoid direct contact with jewellery or metal surfaces as Driclor will cause their discolouration.

BREAST-FEEDING

If you are breast-feeding, Driclor must not get on your breasts to prevent your baby from accidentally swallowing it or being splashed with it. If your breasts are splashed with Driclor, wash the affected areas thoroughly before breast-feeding.

ADVERSE REACTIONS

Very common adverse reactions

They can affect more than 1 in 10 people.

- Irritation of treated skin

Uncommon adverse reactions

They can affect up to 1 in 1,000 people.

The following adverse reactions have occurred on the treated skin:

- Allergic reaction including red, swollen, inflamed skin (dermatitis)
- Pain
- Itching (pruritus)
- Red skin (erythema)
- Rash
- Skin burning sensation

OVERDOSE

Using too much Driclor may cause skin irritation. Wash the skin with plenty of water and ask your pharmacist or doctor for advice. If you accidentally swallow Driclor, you may experience nausea and burning in your mouth. Rinse your mouth thoroughly and ask your pharmacist or doctor for advice.

For external use only. Adults and adolescents aged 12+.

Keep out of the reach of children. Store in temperatures not exceeding 25°C.

Composition: Alcohol, Aluminium Chloride, Aqua

GSK Consumer Healthcare, Levice 934 01, SK

Distributor in the Czech Republic: GlaxoSmithKline Consumer Healthcare Czech Republic s.r.o., Hvězdzdova 1 734/2c, 140 00 Prague 4

Best used by: see packaging. (Exp.)

✉ cz.info@gsk.com
www.gsk.cz

620000
0040140